

Mathematics 554H/701I Homework

Before going on to connectedness, let us practice a bit with compactness. Recall that a metric space is compact if every open cover has a finite subcover.

The following was on the test

Problem 1. Show that every compact metric space can be covered by a finite number of open balls of radius one.

Solution: Let E be a compact metric space. Then the collection $\mathcal{U} = \{B(p, 1) : p \in E\}$ is an open cover of E . (For if $p \in E$, then $p \in B(p, 1) \in \mathcal{U}$.) Because E is compact this open cover will have a finite subcover $\mathcal{U}_0 = \{B(x_1, 1), B(x_2, 1), \dots, B(x_n, 1)\}$. Then \mathcal{U}_0 is the required finite cover of E by open balls of radius 1. \square

We say that a metric space, E , is **totally bounded** if and only if for every positive real number r the space E can be covered by a finite number of balls of radius r .

Proposition 1. *Every compact metric space is totally bounded.*

Problem 2. Prove this. *Hint:* Look at the solution to Problem 1. \square

Problem 3. Let E be a compact metric space. Show that there is a finite subset $F \subseteq E$ such that every point of E is within a distance of $1/100$ of a point of F . \square

We now get back to connectedness.

Definition 2. Let E be a metric space. Then a subset $A \subseteq E$ is **clopen** if and only if it is both open and closed.

Proposition 3. *Let E be a metric space. Then the following are equivalent.*

- (a) E has a clopen subset A with $A \neq \emptyset$ and $A \neq E$.
- (b) There are non-empty disjoint open subsets U and V of E with $E = U \cup V$.
- (c) There are non-empty disjoint closed subsets U and V of E with $E = U \cup V$.

Problem 4. Prove this. *Hint:* See class notes. \square

Definition 4. A metric space E is connected if and only if the only clopen sets in E are E and \emptyset . \square

Definition 5. The sets U and V are a **disconnection** of the metric space E if and only if U and V are open, non-empty, disjoint, and $E = U \cup V$. (In light of Proposition 3 we could also assume that U and V are both closed.) \square

Thus a metric space is connected if and only if it does not have a disconnection. This makes the strategy for showing something is disconnected straightforward: find a disconnection.

Problem 5. Show the following sets are disconnected.

- (a) The rational numbers.
- (b) The set $E = \{(x, y) : x, y \in \mathbb{R} \text{ and } |x| \leq 1, 10 < |y| < 11\}$ *Hint:* Draw a picture. \square

Problem 6. If E is a subset of the real numbers and there are points $a, b \in E$ and a point between a and b that is not in E , then E is not connected. *Hint:* See class notes. \square

Problem 7. Let $r > 0$ and p, q points in a metric space with $d(p, q) > 2r$. Show that the set $E = B(p, r) \cup B(q, r)$ is not connected. \square

Showing sets are connected is more work. Here is a way to reduce showing that a space is connected by showing that it is the union of connected subsets all of which contain the same point.

Problem 8. Let $E = S_1 \cup S_2$ where S_1 and S_2 are both connected and there is a point x_0 that is in both S_1 and S_2 . Then E is connected.

Solution: We need to show that the only clopen subsets of E are E and \emptyset . That is we need to show that if A is a non-empty clopen subset of E , then $A = E$. As $A \subseteq E = S_1 \cup S_2$, then will intersect at least one of S_1 or S_2 . Assume that $A \cap S_1 \neq \emptyset$. Then $A \cap S_1$ is a non-empty clopen subset of S_1 and S_1 is connected, therefore $A \cap S_1 = S_1$. That is $A \subseteq S_1$. As $x_0 \in S_1 \subseteq A$ we have $x_0 \in A$. Therefore $x_0 \in A \cap S_2$ and thus $A \cap S_2 \neq \emptyset$. Then $A \cap S_2$ is a non-empty clopen subset of S_2 and S_2 is connected and therefore $A \cap S_2 = S_2$. Thus $S_2 \subseteq A$. Therefore we have $A \subseteq E = S_1 \cup S_2 \subseteq E$. This shows that $A = E$ which is just what we needed to complete the proof. \square

Proposition 6. Let E be a metric space and $\{S_\alpha\}_{\alpha \in I}$ a collection of connected subsets of E such that for some point $x_0 \in E$ with

$$x_0 \in \bigcap_{\alpha \in I} S_\alpha$$

and

$$E = \bigcup_{\alpha \in I} S_\alpha.$$

Then E is connected.

Problem 9. Prove this. *Hint:* Look at the solution of Problem 8.

Theorem 7. Every interval in \mathbb{R} is connected.

Problem 10. Prove this. *Hint:* Let E be in interval in \mathbb{R} . The property of E that we will use is that if $a, b \in E$ then any real number between a and b is in E . Towards a contradiction assume that E is not connected. Then E is a disjoint union $E = U \cup V$ where U and V are both open and closed in E . Let $a \in U$ and $b \in V$. By possibly changing the names we can assume that $a < b$. Let

$$A = [a, b] \cap U \quad \text{and} \quad B = [a, b] \cap V.$$

and set

$$c = \sup(A)$$

Now show

- (a) A and B are clopen in $[a, b]$.
- (b) $a < c < b$ and $c \in E$ (to start note that as A is open in $[a, b]$ there is a ball $B(a, r)$ with $B(a, r) \cap [a, b] = [a, r) \subseteq A$ and likewise there is a $r' > 0$ such that $(b - r', b] \subseteq B$).
- (c) c is an adherent point of A (this just uses that $c = \sup(A)$)
- (d) c is an adherent point of B . (if not explain why there is $r > 0$ with $(c - r, c + r) \subseteq A$ and why this contradicts that $c = \sup(A)$.)
- (e) As A and B are closed they contain their adherent points and so c is in both A and B .
- (f) Explain why we have a contradiction. □

Problem 11. Let $E \subseteq \mathbb{R}$ such that if $a, b \in E$ and $c \in \mathbb{R}$ is between a and b , then $c \in E$. Show that E is an interval. □