

Mathematics 554 Homework.

Here is a problem to review some of what we know about metric spaces. The reverse triangle inequality is

$$|d(p, x) - d(q, x)| \leq d(p, q).$$

This implies, as we have seen before,

Proposition 1. *If $\lim_{n \rightarrow \infty} p_n = p$ in a metric space, then for any q we have*

$$\lim_{n \rightarrow \infty} d(p_n, q) = d(p, q).$$

Proof. By the reverse triangle inequality

$$|d(p_n, q) - d(p, q)| \leq d(p_n, p).$$

Let $\varepsilon > 0$. Then there is a N so that $n \leq N$ implies $d(p_n, p) < \varepsilon$. But using the reverse triangle inequality just given we have

$$|d(p_n, q) - d(p, q)| \leq d(p_n, p) < \varepsilon$$

which shows $\lim_{n \rightarrow \infty} d(p_n, q) = d(p, q)$. \square

Remark 2. Another way to think of this is that the reverse triangle inequality gives that $f(x) := d(x, q)$ is a Lipschitz function with Lipschitz constant $M = 1$. And we have shown that for a Lipschitz function that $\lim_{n \rightarrow \infty} p_n = p$ implies $\lim_{n \rightarrow \infty} f(p_n) = f(p)$. \square

Proposition 3. *If $\lim_{n \rightarrow \infty} p_n = p$ and $\lim_{n \rightarrow \infty} q_n = q$ in a metric space, then*

$$\lim_{n \rightarrow \infty} d(p_n, q_n) = d(p, q).$$

Problem 1. Prove this. *Hint:* One one to start to justify the following:

$$\begin{aligned} |d(p_n, q_n) - d(p, q)| &= |d(p_n, q_n) - d(p, q_n) + d(p, q_n) - d(p, q)| \\ &\leq |d(p_n, q_n) - d(p, q_n)| + |d(p, q_n) - d(p, q)| \\ &\leq d(p_n, p) + d(q_n, q) \end{aligned}$$

and do an $\varepsilon/2$ proof.

Here are some applications showing one of the many ways compactness is used.

Let S be a set in a metric space (or just in the plane, \mathbb{R}^2 , if you like). Then the **diameter** of S is

$$\text{Diameter}(S) := \sup\{d(p, q) : p, q \in S\}.$$

That is the diameter is the least upper bound on the distances between points in S . We would like to say that it is the greatest possible distance between point of S this is not always the case. For example if $S = (0, 1)$, an open interval in \mathbb{R} , then $\text{Diameter}(S) = 1$, but if $p, q \in (0, 1) = S$ we have $d(p, q) = |p - q| < 1$. On the other hand if S is the closed interval $[0, 1]$, then

the maximum distance is realised, as $0, 1 \in [0, 1]$ and $\text{Diameter}([0, 1]) = 1 = |1 - 0|$.

In general we can find points that are at a maximal distance apart in a set when the set is compact.

Proposition 4. *Let S be a compact set in a metric space. Then there are points $p_0, q_0 \in S$ with*

$$\text{Diameter} = d(p_0, q_0).$$

Proof. We have proven that compact sets in metric spaces are bounded, so the diameter is finite. Let $D = \text{Diameter}(S)$. By the definition of Diameter as $\sup(\{d(p, q) : p, q \in S\})$ we have for each $n \in \mathbb{N}$ a pair p_n, q_n such that

$$D - \frac{1}{n} < d(p_n, q_n) \leq D.$$

This implies $|D - d(p_n, q_n)| < 1/n$ and therefore

$$\lim_{n \rightarrow \infty} d(p_n, q_n) = D.$$

(For, as we have done before, if $\varepsilon > 0$ and $N = 1/\varepsilon$, then $n > N$ implies $|D - d(p_n, q_n)| < 1/n < \varepsilon$.)

$$\lim_{n \rightarrow \infty} d(p_n, q_n) = D.$$

As S is compact, it is also sequentially compact. Thus there is a subsequence $\langle p_{n_k} \rangle_{k=1}^\infty$ so that

$$\lim_{k \rightarrow \infty} p_{n_k} = p_0$$

for some $p \in S$. Now we have to do the subsequence of a subsequence trick. There is a subsequence $\langle q_{n_{k_j}} \rangle_{j=1}^\infty$ of $\langle q_{n_k} \rangle_{k=1}^\infty$ so that

$$\lim_{j \rightarrow \infty} q_{n_{k_j}} = q_0$$

for some q_0 in S . Using that subsequences of convergent sequences converge to the same limit as the ordinal sequence together with Proposition 3 gives

$$d(p_0, q_0) = \lim_{j \rightarrow \infty} d(p_{n_{k_j}}, q_{n_{k_j}}) = D.$$

Thus p_0 and q_0 satisfy the conclusion of the proposition. \square

Proposition 5. *Let A and B be compact subsets of the metric space and let*

$$M = \inf(\{d(a, b) : a \in A, \text{ and } b \in B\}).$$

Then there are $a_0 \in A$ and $b_0 \in B$ so that

$$d(a_0, b_0) = M.$$

Problem 2. Prove this. *Hint:* This is very much like the proof of Proposition 4. By the definition of M as an infimum for each $n \in \mathbb{N}$ there are $a_n \in A$ and $b_n \in B$ such that

$$M \leq d(a_n, b_n) < M + \frac{1}{n}$$

and this implies $\lim_{n \rightarrow \infty} d(a_n, b_n) = M$. Now use sequential compactness and a subsequence of a subsequence argument to get the existence of a_0 and b_0 so that $d(a_0, b_0) = M$. \square

Problem 3. Let A be a subset of $[a, b]$ with $a \in A$ and so that A is both open and closed in $[a, b]$. Show $A = [a, b]$. *Hint:* Toward a contradiction assume that $A \neq [a, b]$. Let $B = [a, b] \setminus A$ be the complement of A in $[a, b]$. Explain why B is also both open and closed in $[a, b]$. Let $\beta = \inf(B)$ and explain why $\beta \in B$. Then explain why β is an adherent point of A and why this implies $\beta \in A$. We now have that $\beta \in A \cap B$, which contradicts that B is the complement of A . \square

Problem 4. Prove that any compact metric space, E , is complete. *Hint:* You need to show that any Cauchy sequence in E converges. So you probably want to use that compact spaces are sequentially compact. \square

The next problem is off of one of my previous Math 554 exams. (It was during pandemic and thus was take home so time was not an issue.)

Problem 5. Let define a sequence of real numbers $\langle x_n \rangle_{n=1}^{\infty}$ by

$$\begin{aligned} x_1 &= 1 \\ x_2 &= \frac{3}{4}x_1 + 12 \\ x_3 &= \frac{3}{4}x_2 + 12 \\ x_4 &= \frac{3}{4}x_3 + 12 \end{aligned}$$

and in general

$$x_n = \frac{3}{4}x_{n-1} + 12.$$

- (a) Compute x_2, x_3, x_4 .
- (b) Use induction to show the sequence is increasing.
- (c) Use induction to show $x_n \leq 100$ for all n .
- (d) Prove $\langle x_n \rangle_{n=1}^{\infty}$ converges and find its limit. \square

In the class where the previous problem was given the Banach Fixed Point Theorem was not covered. Using the fixed point theorem we can generalize the problem.

Problem 6. Recall that a map $f: E \rightarrow E$ from a metric space to itself is **contraction** if and only if there is a constant ρ with $0 \leq \rho < 1$ such that

$$d(f(p), f(q)) \leq \rho d(p, q)$$

for all $p, q \in E$. The Banach Fixed Point Theorem is that if f is a contraction and the space E is complete, then f has a unique fixed point (that is a point

p with $f(p) = p$ and that this fixed point is the limit of an sequence defined by choosing any point $p_0 \in E$ and defining p_n for $n \geq 1$ by

$$x_n = f(x_{n-1}).$$

Then the unique is the limit

$$p_* = \lim_{n \rightarrow \infty} p_n.$$

Now let $a, b \in \mathbb{R}$ be constants with $|a| < 1$ and define $f: \mathbb{R} \rightarrow \mathbb{R}$ by

$$f(x) = ax + b.$$

- (a) Show f is a contraction.
- (b) Therefore the Banach Fixed Point Theorem implies that if x_0 is any real number and for $n \geq 1$

$$x_n = f(x_{n-1}) = ax_{n-1} + b$$

that this sequence converges to the unique fixed point of f . Find the fixed point and in the case where $a = 3/4$ and $b = 12$ compare with the answer to Problem 5. \square

Problem 7. (Anther problem off of an old exam.) Let S be the subset of \mathbb{R}^2 defined by

$$S = \{(x, y) : 0 < x^2 + y^2 \leq 1\}.$$

- (a) Draw a picture of S .
- (b) Show that S is not sequentially compact. \square

Problem 8. This is anther review problem. Let $f: E \rightarrow \mathbb{R}$ be Lipschitz, say $|f(p) - f(q)| \leq Md(p, q)$ for all $p, q \in E$. Then for any $c \in \mathbb{R}$ the sets

$$f^{-1}((-\infty, c)) = \{p : f(p) < c\}, \quad \text{and} \quad f^{-1}((c, \infty)) = \{p : f(p) > c\}$$

are both open. Choose one these two and prove it is open. (No extra points for doing both.) \square

Proposition 6. Let E be a connected metric space and $f: E \rightarrow \mathbb{R}$ a Lipschitz function. Assume there are points $p_1, p_2 \in E$ with $f(p_1) < 0$ and $f(p_2) > 0$. Then there is a point $q \in E$ with $f(q) = 0$.

Problem 9. Prove this. *Hint:* Assume that it is false. Then use Problem 8 to show that

$$A := \{p : f(p) < 0\} \quad \text{and} \quad B := \{p : f(p) > 0\}$$

is a disconnection of E , contradicting that E is connected.

Problem 10. Anther problem off of an old exam. Let $\langle x_n \rangle_{n=1}^\infty$ be a sequence of real numbers with $\lim_{n \rightarrow \infty} x_n = L$. Give a ε, N proof that $\lim_{n \rightarrow \infty} x_n^2 = L^2$.