

Mathematics 552 Homework, March 9, 2020

In class on Friday we showed

Theorem 1. *Let $f(z)$ be a complex valued function defined on a curve γ . Assume there is a constant M such that*

$$|f(z)| \leq M$$

for all z on the curve γ . Then

$$\left| \int_{\gamma} f(z) dz \right| \leq ML(\gamma)$$

where $L(\gamma)$ is the length of γ . □

Here is an example of the use of this result. Let Assume that $|f(z)| \leq 5$ on the circle $|z - 2i| = 3$. Then

$$\begin{aligned} \left| \int_{|z-2i|=3} f(z) dz \right| &\leq 5 \times \text{Length of circle radius 3} \\ &= 5 \times 2\pi(3) \\ &= 30\pi. \end{aligned}$$

We have also shown that if $f(z)$ is on an inside of a simple closed curve γ that for a inside of γ that the derivative $f'(a)$ is given by

$$f'(a) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{(z-a)^2} dz.$$

Proposition 2. *Let $f(z)$ be analytic on and inside of the circle $|z - a| = r$ and assume*

$$|f(z)| \leq M \quad \text{on the circle } |z - a| = r.$$

Then

$$|f'(a)| \leq \frac{M}{r}.$$

Problem 1. Prove this along the following lines.

(a) Explain why for z on the circle $|z - a| = r$ the equality

$$\left| \frac{1}{(z-a)^2} \right| = \frac{1}{r^2}$$

holds.

(b) Let $F(z) = \frac{f(z)}{(z-a)^2}$ and use $|f(z)| \leq M$ and part (a) to show that on the circle $|z - a| = r$ the inequality

$$|F(z)| \leq \frac{M}{r^2}.$$

(c) Use Theorem 1 to explain why

$$\left| \int_{|z-a|=r} F(z) dz \right| \leq \frac{M}{r^2} \times 2\pi r = \frac{2\pi M}{r}.$$

(d) With this notation note and the formula for $f'(a)$ we have

$$|f'(a)| = \left| \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{(z-a)^2} dz \right| = \left| \frac{1}{2\pi i} \int_{|z-a|=r} F(z) dz \right|.$$

Use this to complete the proof of Proposition 2. \square

Definition 3. A function $f(z)$ that is analytic on all of \mathbb{C} is called an *entire function*.

The following is one of the more famous results in complex analysis.

Theorem 4 (Louisville's Theorem). *A bounded entire function is constant.*

Problem 2. Prove this. *Hint:* To show a function is constant it is enough to show that its derivative is always zero. That is we want to show $f'(a) = 0$ for all a . What we know is that $f(z)$ is bounded. Explicitly this means there is a constant M so that $|f(z)| \leq M$ for all z . Let $r > 0$ and explain by

$$|f'(a)| \leq \frac{M}{r}.$$

Now take the limit as $r \rightarrow \infty$ to conclude that $|f'(a)| = 0$, and so $f'(a) = 0$ which finishes the proof.

One of the important applications of Louisville's Theorem is showing that every polynomial with complex coefficients has a root.

We know the *triangle inequality* for complex numbers

$$|z + w| \leq |z| + |w|.$$

Problem 3. Use the triangle inequality to show for any complex numbers a, b that

$$|a + b| \geq |a| - |b|.$$

Hint: In the triangle inequality let $z = a + b$ and $w = -b$.

Problem 4. Use the last problem repeatedly to show

$$|a + b_1 + b_2 + \cdots + b_n| \geq |a| - |b_1| - |b_2| - \cdots - |b_n|.$$

Instead of working with polynomials of degree n , it will simplify notation if we work with polynomials of degree 3. All the basic ideas are the same.

Problem 5. Let $p(z) = z^3 + b_2 z^2 + b_1 z + b_0$. Show

$$|p(z)| \geq |z|^3 \left(1 - \frac{|b_2|}{|z|} - \frac{|b_1|}{|z|^2} - \frac{|b_0|}{|z|^3} \right).$$

Problem 6. With notation as in Problem 5 show that if $R = \max\{1, 6|b_2|, 6|b_1|, 6|b_0|\}$ then show that for $|z| \geq R$ (that is $|z| \geq 1$, $|z| \geq 6|b_2|$, $|z| \geq 6|b_1|$) that the following hold

(a) $\frac{1}{|z|^3} \leq \frac{1}{|z|^2} \leq \frac{1}{|z|} \leq 1$. *Hint:* This only uses $|z| \geq 1$.

(b) $\frac{|b_2|}{|z|} \leq \frac{1}{6}$. *Hint:* This uses $|z| \geq 6|b_2|$.

(c) $\frac{|b_1|}{|z|^2} \leq \frac{1}{6}$. *Hint:* This uses $|z| \geq 6|b_1|$ and part (a).

(d) $\frac{|b_0|}{|z|^3} \leq \frac{1}{6}$. *Hint:* This uses $|z| \geq 6|b_0|$ and part (a).

(e) $|p(z)| \geq \frac{|z|^3}{2} \geq \frac{1}{2}$. *Hint:* This uses parts (b), (c), (d) and Problem 3.

Theorem 5 (Fundamental Theorem of Algebra). *Let $p(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0$ be a complex polynomial of degree $n \geq 1$. Then $p(z)$ has at least one complex root. That is there is at least one complex number r with $p(r) = 0$.*

The following problems will give a proof in the case of $n = 3$. The general case is not much harder. So we start with the polynomial

$$p(z) = a_3 z^3 + a_2 z^2 + a_1 z + a_0$$

with $a_3 \neq 0$.

To start we note that by dividing by a_n we have that solving

$$p(z) = a_3 z^3 + a_2 z^2 + a_1 z + a_0 = 0$$

is the same as solving

$$z^3 + \frac{a_2}{a_3} z^2 + \frac{a_1}{a_3} z + \frac{a_0}{a_3} = 0$$

so there is no loss of generality in assuming that the lead coefficient of $p(z)$ is one. That is $p(z)$ is of the form

$$p(z) = z^3 + b_2 z^2 + b_1 z + b_0.$$

Assume, towards a contradiction, that $p(z)$ has no roots. That is $p(z) \neq 0$ for all $z \in \mathbb{C}$. Define a new function $f(z)$ by

$$f(z) = \frac{1}{p(z)}.$$

Problem 7. Explain why $f(z)$ is an entire function. That is explain why $f(z)$ is differentiable at all points.

Problem 8. Let R be as in Problem 6. Show

$$|z| \geq R \quad \text{implies} \quad |f(z)| \leq 2.$$

Problem 9. The function $|f(z)|$ is continuous on the closed bounded set $\{z : |z| \leq R\}$, so there is a constant C such that

$$|z| \leq R \quad \text{implies} \quad |f(z)| \leq C.$$

(This is a basic fact from Mathematics 554, so you don't have to prove it, just copy it down to get credit.)

Problem 10. Let R be as in Problem 6 and set $M = \max\{2, C\}$. Combine Problems 8 and 9 to show

$$|f(z)| \leq M$$

for all $z \in \mathbb{C}$.

Problem 11. Now show that $f(z) = \frac{1}{p(z)}$ is constant and therefore $p(z)$ is also constant.

And finally

Problem 12. To finish the proof explain why the assumption $p(z)$ has no roots leads to a contradiction.